

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number  
WO 01/49281 A2

- (51) International Patent Classification<sup>7</sup>: A61K 31/00
- (21) International Application Number: PCT/US00/35715
- (22) International Filing Date:  
28 December 2000 (28.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/173,958 30 December 1999 (30.12.1999) US  
09/748,748 26 December 2000 (26.12.2000) US
- (71) Applicant: PROTEO TECH, INC. [US/US]; 12026  
115th Avenue N.E., Kirkland, WA 98034 (US).
- (72) Inventors: CASTILLO, Gerardo, M.; 11725-82nd Ave.  
South, Seattle, WA 98178 (US). CHOI, Paula, Y.; Apt.  
108, 11119 Woodenville Drive, Bothell, WA 98011 (US).  
SNOW, Alan, D.; 3812 167th Pl. S.W., Lynnwood, WA  
98037 (US).
- (74) Agents: KURZ, Walter et al.; Heller Ehrman White &  
McAuliffe LLP, 275 Middlefield Road, Menlo Park, CA  
94025-3506 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AT  
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility  
model), DK, DK (utility model), DM, DZ, EE, EE (utility  
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK  
(utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— Without international search report and to be republished  
upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



WO 01/49281 A2

(54) Title: POLYHYDROXYLATED AROMATIC COMPOUNDS FOR THE TREATMENT OF AMYLOIDOSIS AND  $\alpha$ -SYNU-  
CLEIN FIBRIL DISEASES

(57) Abstract: Polyhydroxylated aromatic compounds, and compositions containing them, are useful for the treatment of amyloido-  
sis, especially Alzheimer's disease, and for the treatment of diseases characterized by  $\alpha$ -synuclein fibril formation, especially Lewy  
body disease and Parkinson's disease.

**POLYHYDROXYLATED AROMATIC COMPOUNDS FOR THE TREATMENT  
OF AMYLOIDOSIS AND  $\alpha$ -SYNUCLEIN FIBRIL DISEASES**

5                                   **BACKGROUND OF THE INVENTION**

**FIELD OF THE INVENTION**

This invention relates to the use of certain polyhydroxylated aromatic compounds, and compositions containing them, for the treatment of amyloidosis, especially Alzheimer's disease, and the treatment of diseases characterized by  $\alpha$ -synuclein fibril  
10 formation, especially Lewy body disease and Parkinson's disease.

**DESCRIPTION OF THE RELATED ART**

**Amyloid and amyloidosis**

Amyloid is a generic term referring to a group of diverse but specific extracellular  
15 protein deposits which all have common morphological properties, staining characteristics, and X-ray diffraction spectra. Regardless of the nature of the amyloid protein deposited all amyloids have the following characteristics: 1) showing an amorphous appearance at the light microscopic level, appearing eosinophilic using hematoxylin and eosin stains; 2) staining with Congo red and demonstrating a red/green  
20 birefringence as viewed under polarized light (Puchtler et al., *J. Histochem. Cytochem.* 10:355-364, 1962), 3) containing a predominant beta-pleated sheet secondary structure, and 4) ultrastructurally consisting of non-branching fibrils of indefinite length and with a diameter of 7-10 nm.

25           Amyloidoses today are classified according to the specific amyloid protein deposited. The amyloids include, but are not limited to, the amyloid associated with Alzheimer's disease, Down's syndrome and hereditary cerebral hemorrhage with amyloidosis of the Dutch type (where the specific amyloid is referred to as beta-amyloid protein or A $\beta$ ), the amyloid associated with chronic inflammation, various forms of  
30 malignancy and familial Mediterranean fever (where the specific amyloid is referred to as AA amyloid or inflammation-associated amyloid), the amyloid associated with multiple myeloma and other B-cell dyscrasias (where the specific amyloid is referred to as AL

amyloid), the amyloid associated with type II diabetes (where the specific amyloid is referred to as amylin or islet amyloid), the amyloid associated with the prion diseases including Creutzfeldt-Jakob disease, Gerstmann-Straussler syndrome, kuru, and scrapie (where the specific amyloid is referred to as PrP amyloid), the amyloid associated with  
5 long-term hemodialysis and carpal tunnel syndrome (where the specific amyloid is referred to as beta<sub>2</sub>-microglobulin amyloid), the amyloid associated with senile cardiac amyloid and familial amyloidotic polyneuropathy (where the specific amyloid is referred to as prealbumin or transthyretin amyloid), and the amyloid associated with endocrine tumors such as medullary carcinoma of the thyroid (where the specific amyloid is  
10 referred to as variants of procalcitonin).

Although amyloid deposits in clinical conditions share common physical properties relating to the presence of a beta-pleated sheet conformation, it is now clear that many different chemical types exist and additional ones are likely to be described in the future. It is currently thought that there are several common pathogenetic mechanisms  
15 that may be operating in amyloidosis in general. In many cases, a circulating precursor protein may result from overproduction of either intact or aberrant molecules (for example, in plasma cell dyscrasias), reduced degradation or excretion (serum amyloid A in some secondary amyloid syndromes and beta<sub>2</sub>-microglobulin in long-term hemodialysis), or genetic abnormalities associated with variant proteins (for example,  
20 familial amyloidotic polyneuropathy). Proteolysis of a larger protein precursor molecule occurs in many types of amyloidosis, resulting in the production of lower molecular weight fragments that polymerize and assume a beta-pleated sheet conformation as tissue deposits, usually in an extracellular location. The precise mechanisms involved and the aberrant causes leading to changes in proteolytic processing and/or translational  
25 modification are not known in most amyloids.

Systemic amyloids which include the amyloid associated with chronic inflammation, various forms of malignancy and familial Mediterranean fever (i.e. AA amyloid or inflammation-associated amyloidosis) (Benson and Cohen, *Arth. Rheum.* 22:36-42, 1979; Kamei et al, *Acta Path. Jpn.* 32:123-133, 1982; McAdam et al., *Lancet*  
30 2:572-573, 1975; Metaxas, *Kidney Int.* 20:676-685, 1981), and the amyloid associated with multiple myeloma and other B-cell dyscrasias (i.e. AL amyloid) (Harada et al., *J.*

*Histochem. Cytochem.* 19:1-15, 1971), as examples, are known to involve amyloid deposition in a variety of different organs and tissues generally lying outside the central nervous system. Amyloid deposition in these diseases may occur, for example, in liver, heart, spleen, gastrointestinal tract, kidney, skin, and/or lungs (Johnson et al, *N. Engl. J. Med.* 321:513-518, 1989). For most of these amyloidoses, there is no apparent cure or effective treatment and the consequences of amyloid deposition can be detrimental to the patient. For example, amyloid deposition in the kidney may lead to renal failure, whereas amyloid deposition in the heart may lead to heart failure. For these patients, amyloid accumulation in systemic organs leads to eventual death generally within 3-5 years. Other amyloidoses may affect a single organ or tissue such as observed with the A $\beta$  amyloid deposits found in the brains of patients with Alzheimer's disease and Down's syndrome: the PrP amyloid deposits found in the brains of patients with Creutzfeldt-Jakob disease, Gerstmann-Straussler syndrome, and kuru; the islet amyloid (amylin) deposits found in the islets of Langerhans in the pancreas of 90% of patients with type II diabetes (Johnson et al, *N. Engl. J. Med.* 321:513-518, 1989; *Lab. Invest.* 66:522-535, 1992); the beta<sub>2</sub>-microglobulin amyloid deposits in the medial nerve leading to carpal tunnel syndrome as observed in patients undergoing long-term hemodialysis (Geyjo et al, *Biochem. Biophys. Res. Comm.* 129:701-706, 1985; *Kidney Int.* 30:385-390, 1986); the prealbumin/transthyretin amyloid observed in the hearts of patients with senile cardiac amyloid; and the prealbumin/transthyretin amyloid observed in peripheral nerves of patients who have familial amyloidotic polyneuropathy (Skinner and Cohen, *Biochem. Biophys. Res. Comm.* 99:1326-1332, 1981; Saraiva et al, *J. Lab. Clin. Med.* 102:590-603, 1983; *J. Clin. Invest.* 74:104-119, 1984; Tawara et al, *J. Lab. Clin. Med.* 98:811-822, 1989).

25

#### **Alzheimer's disease and the aging population**

Alzheimer's disease is a leading cause of dementia in the elderly, affecting 5-10% of the population over the age of 65 years (*A Guide to Understanding Alzheimer's Disease and Related Disorders*, Jorm, ed., New York University Press, New York, 1987). In Alzheimer's disease, the parts of the brain essential for cognitive processes such as memory, attention, language, and reasoning degenerate, robbing victims of much that

30

makes us human, including independence. In some inherited forms of Alzheimer's disease, onset is in middle age, but more commonly, symptoms appear from the mid-60's onward. Alzheimer's disease today affects 4-5 million Americans, with slightly more than half of these people receiving care at home, while the others are in many different health care institutions. The prevalence of Alzheimer's disease and other dementias doubles every 5 years beyond the age of 65, and recent studies indicate that nearly 50% of all people age 85 and older have symptoms of Alzheimer's disease (*1999 Progress Report on Alzheimer's Disease*, National Institute on Aging/National Institute of Health). 13% (33 million people) of the total population of the United States are age 65 and older, and this percentage will climb to 20% by the year 2025 (*1999 Progress Report on Alzheimer's Disease*).

Alzheimer's disease also puts a heavy economic burden on society. A recent study estimated that the cost of caring for one Alzheimer's disease patient with severe cognitive impairments at home or in a nursing home, is more than \$47,000 per year (*A Guide to Understanding Alzheimer's Disease and Related Disorders*). For a disease that can span from 2 to 20 years, the overall cost of Alzheimer's disease to families and to society is staggering. The annual economic toll of Alzheimer's disease in the United States in terms of health care expenses and lost wages of both patients and their caregivers is estimated at \$80 to \$100 billion (*1999 Progress Report on Alzheimer's Disease*).

Tacrine hydrochloride ("Cognex"), the first FDA approved drug for Alzheimer's disease, is a acetylcholinesterase inhibitor (Cutler and Sramek, *N. Engl. J. Med.* 328:808-810, 1993). However, this drug has showed limited success in producing cognitive improvement in Alzheimer's disease patients and initially had major side effects such as liver toxicity. The second more recently FDA approved drug, donepezil ("Aricept"), which is also an acetylcholinesterase inhibitor, is more effective than tacrine, by demonstrating slight cognitive improvement in Alzheimer's disease patients (Barner and Gray; *Ann. Pharmacotherapy* 32:70-77, 1998; Rogers and Friedhoff, *Eur. Neuropsych.* 8:67-75, 1998), but is not believed to be a cure. Therefore, it is clear that there is a need for more effective treatments for Alzheimer's disease patients.

### Amyloid as a therapeutic target for Alzheimer's disease

Alzheimer's disease is characterized by the deposition and accumulation of a 39-43 amino acid peptide termed the beta-amyloid protein, A $\beta$  or  $\beta$ /A4 (Glennner and Wong, *Biochem. Biophys. Res. Comm.* 120:885-890, 1984; Masters et al., *Proc. Natl. Acad. Sci. USA* 82:4245-4249, 1985; Husby et al., *Bull. WHO* 71:105-108, 1993). A $\beta$  is derived by protease cleavage from larger precursor proteins termed beta-amyloid precursor proteins (or  $\beta$ PPs) of which there are several alternatively spliced variants. The most abundant forms of the  $\beta$ PPs include proteins consisting of 695, 751 and 770 amino acids (Tanzi et al., *Nature* 331:528-530, 1988; Kitaguchi et al., *Nature* 331:530-532, 1988; Ponte et al., *Nature* 331:525-527, 1988).

The small A $\beta$  peptide is a major component which makes up the amyloid deposits of "plaques" in the brains of patients with Alzheimer's disease. In addition, Alzheimer's disease is characterized by the presence of numerous neurofibrillary "tangles", consisting of paired helical filaments which abnormally accumulate in the neuronal cytoplasm (Grundke-Iqbal et al., *Proc. Natl. Acad. Sci. USA* 83:4913-4917, 1986; Kosik et al., *Proc. Natl. Acad. Sci. USA* 83:4044-4048, 1986; Lee et al., *Science* 251:675-678, 1991). The pathological hallmark of Alzheimer's disease is therefore the presence of "plaques" and "tangles", with amyloid being deposited in the central core of the plaques. The other major type of lesion found in the Alzheimer's disease brain is the accumulation of amyloid in the walls of blood vessels, both within the brain parenchyma and in the walls of meningeal vessels which lie outside the brain. The amyloid deposits localized to the walls of blood vessels are referred to as cerebrovascular amyloid or congophilic angiopathy (Mandybur, *J. Neuropath. Exp. Neurol.* 45:79-90, 1986; Pardridge et al., *J. Neurochem.* 49:1394-1401, 1987).

For many years there has been an ongoing scientific debate as to the importance of "amyloid" in Alzheimer's disease, and whether the "plaques" and "tangles" characteristic of this disease were a cause or merely a consequence of the disease. Within the last few years, studies now indicate that amyloid is indeed a causative factor for Alzheimer's disease and should not be regarded as merely an innocent bystander. The Alzheimer's A $\beta$  protein in cell culture has been shown to cause degeneration of nerve cells within short periods of time (Pike et al., *Br. Res.* 563:311-314, 1991; *J. Neurochem.*

64:253-265, 1995). Studies suggest that it is the fibrillar structure (consisting of a predominant beta-pleated sheet secondary structure), characteristic of all amyloids, that is responsible for the neurotoxic effects. A $\beta$  has also been found to be neurotoxic in slice cultures of hippocampus (Harrigan et al., *Neurobiol. Aging* 16:779-789, 1995) and induces nerve cell death in transgenic mice (Games et al., *Nature* 373:523-527, 1995; Hsiao et al., *Science* 274:99-102, 1996). Injection of the Alzheimer's A $\beta$  into rat brain also causes memory impairment and neuronal dysfunction (Flood et al., *Proc. Natl. Acad. Sci. USA* 88:3363-3366, 1991; *Br. Res.* 663:271-276, 1994).

Probably, the most convincing evidence that A $\beta$  amyloid is directly involved in the pathogenesis of Alzheimer's disease comes from genetic studies. It has been discovered that the production of A $\beta$  can result from mutations in the gene encoding, its precursor, beta amyloid precursor protein (Van Broeckhoven et al., *Science* 248:1120-1122, 1990; Murrell et al., *Science* 254:97-99, 1991; Haass et al., *Nature Med.* 1:1291-1296, 1995). The identification of mutations in the beta-amyloid precursor protein gene which causes early onset familial Alzheimer's disease is the strongest argument that amyloid is central to the pathogenetic process underlying this disease. Four reported disease-causing mutations have now been discovered which demonstrate the importance of A $\beta$  in causing familial Alzheimer's disease (reviewed in Hardy, *Nature Genet.* 1:233-234, 1992). All of these studies suggest that providing a drug to reduce, eliminate or prevent fibrillar A $\beta$  formation, deposition, accumulation and/or persistence in the brains of human patients will serve as an effective therapeutic.

Discovery and identification of new compounds or agents as potential therapeutic agents to arrest amyloid deposition, accumulation and/or persistence that occurs in Alzheimer's disease and other amyloidoses are desperately sought.

#### **Parkinson's Disease and $\alpha$ -Synuclein Fibril Formation**

Parkinson's disease is a neurodegenerative disorder that is pathologically characterized by the presence of intracytoplasmic Lewy bodies (Lewy in *Handbuch der Neurologie*, M. Lewandowski, ed., Springer, Berlin, pp. 920-933, 1912; Pollanen et al., *J. Neuropath. Exp. Neurol.* 52:183-191, 1993), the major components of which are filaments consisting of  $\alpha$ -synuclein (Spillantini et al., *Proc. Natl. Acad. Sci. USA*

95:6469-6473, 1998; Arai et al., *Neurosc. Lett.* 259:83-86, 1999), an 140-amino acid protein (Ueda et al., *Proc. Natl. Acad. Sci. USA* 90:11282-11286, 1993). Two dominant mutations in  $\alpha$ -synuclein causing familial early onset Parkinson's disease have been described suggesting that Lewy bodies contribute mechanistically to the degeneration of neurons in Parkinson's disease (Polymeropoulos et al., *Science* 276:2045-2047, 1997; Kruger et al., *Nature Genet.* 18:106-108, 1998). Recently, *in vitro* studies have demonstrated that recombinant  $\alpha$ -synuclein can indeed form Lewy body-like fibrils (Conway et al., *Nature Med.* 4:1318-1320, 1998; Hashimoto et al., *Brain Res.* 799:301-306, 1998; Nahri et al., *J. Biol. Chem.* 274:9843-9846, 1999). Most importantly, both Parkinson's disease-linked  $\alpha$ -synuclein mutations accelerate this aggregation process which suggests that such *in vitro* studies may have relevance for Parkinson's disease pathogenesis.  $\alpha$ -Synuclein aggregation and fibril formation fulfills of the criteria of a nucleation-dependent polymerization process (Wood et al., *J. Biol. Chem.* 274:19509-19512, 1999). In this regard  $\alpha$ -synuclein fibril formation resembles that of Alzheimer's beta-amyloid protein (A $\beta$ ) fibrils.  $\alpha$ -Synuclein recombinant protein, and non-amyloid component (known as NAC-P), which is a 35-amino acid peptide fragment of  $\alpha$ -synuclein, both have the ability to form fibrils when incubated at 37°C, and are positive with amyloid stains such as Congo red (demonstrating a red/green birefringence when viewed under polarized light) and Thioflavin S (demonstrating positive fluorescence) (Hashimoto et al., *Brain Res.* 799:301-306, 1998; Ueda et al., *Proc. Natl. Acad. Sci. USA* 90:11282-11286, 1993).


Parkinson's disease  $\alpha$ -synuclein fibrils, like the A $\beta$  fibrils of Alzheimer's disease, also consist of a predominant beta-pleated sheet structure. We believe, therefore, that compounds found to inhibit Alzheimer's disease A $\beta$  amyloid fibril formation can also be anticipated to be effective in the inhibition of  $\alpha$ -synuclein fibril formation. These compounds would therefore also serve as therapeutics for Parkinson's disease, in addition to having efficacy as a therapeutic for Alzheimer's disease and other amyloid disorders.

The disclosures of these and other documents referred to throughout this application are incorporated herein by reference.




## SUMMARY OF THE INVENTION


In a first aspect, this invention provides a method of treating amyloidosis in a mammal suffering therefrom, comprising administration to the mammal of a therapeutically effective amount of an isolated pure compound selected from the group consisting of the compounds of formula A, formula B, formula C, formula D, and formula E:




Formula A




Formula B



Formula C



Formula D



Formula E

10 where:

R is selected from the group consisting of hydrogen, 2,3-dihydroxybenzoyl, 3,4-dihydroxybenzoyl, 2,3,4-trihydroxybenzoyl, and 3,4,5-trihydroxybenzoyl;

R' is hydrogen or OH;

R<sub>1</sub> and R<sub>2</sub> are independently selected from hydrogen and non-interfering substituents;

15 X is selected from hydrogen and the group consisting of

(a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,

- (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
- 5 (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents,
- (d) sugars, optionally substituted with one or more anionic groups selected from sulfate, phosphate, phosphonate, carboxylate, and sulfonate groups,
- (e) peptides and peptide derivatives, and
- 10 (f) -C(O)R<sub>3</sub> and -C(O)OR<sub>3</sub> (where R<sub>3</sub> is selected from the group consisting of (a) through (e) above); and

Y is hydrogen, hydroxy, C<sub>1-6</sub> alkoxy, benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl), or —OSO<sub>2</sub>R<sub>4</sub> (where R<sub>4</sub> is C<sub>1-6</sub> alkyl or phenyl optionally substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl);

15

and the group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallol, anthralin, anthrarobin, antharufin, apigenin, apigetrin, apiose, baicalein, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose,

20 centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, chrysamminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromyctin, collinomycin, curvularin, cyanidin, cyanidin 3-glucoside, cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside, cyanidin 3-sophoroside, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin,

25 diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscine, fustin, galangin, gallein, gallocyanine, gardenin A, gardenin B, gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecycline, hematein,

30 hydroxysophorobioside, hydroxysophoricoside, icariin, isoquercitrin, kaempferol, kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D,

leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin,  
oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin,  
purpurogallin, quercetagenin, quercimritrin, quinalizarin, quinic acid,  
resistomycin, rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B,  
5 robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone,  
tiron, troxerutin, and tunichrome B1,

and the pharmaceutically acceptable salts thereof.


In preferred embodiments of this first aspect, only one such compound is  
administered; the mammal is a human; and the amyloidosis is selected from the group  
10 consisting of Alzheimer's disease, Down's syndrome, hereditary cerebral hemorrhage  
with amyloidosis of the Dutch type, the amyloidosis of chronic inflammation, the  
amyloidosis of malignancy, familial Mediterranean fever, multiple myeloma, B-cell  
dyscrasias, type II diabetes, the prion diseases, Creutzfeldt-Jakob disease, Gerstmann-  
Straussler syndrome, kuru, scrapie, the amyloidosis associated with long-term  
15 hemodialysis, the amyloidosis associated with carpal tunnel syndrome, senile cardiac  
amyloidosis, familial amyloidotic polyneuropathy, and the amyloidosis associated with  
endocrine tumors, and especially is Alzheimer's disease.

In a second aspect, this invention provides a drug product for the treatment of  
amyloidosis in a mammal suffering therefrom, comprising a container labeled or  
20 accompanied by a label indicating that the drug product is for the treatment of  
amyloidosis, the container containing one or more dosage units each comprising at least  
one pharmaceutically acceptable excipient and, as an active ingredient, an isolated pure  
compound selected from those used in the method of the first aspect of this invention.


In preferred embodiments of this second aspect, the drug product contains only  
25 one such compound, the mammal is a human; and the amyloidosis is selected from the  
group consisting of Alzheimer's disease, Down's syndrome, hereditary cerebral  
hemorrhage with amyloidosis of the Dutch type, the amyloidosis of chronic  
inflammation, the amyloidosis of malignancy, familial Mediterranean fever, multiple  
myeloma, B-cell dyscrasias, type II diabetes, the prion diseases, Creutzfeldt-Jakob  
30 disease, Gerstmann-Straussler syndrome, kuru, scrapie, the amyloidosis associated with  
long-term hemodialysis, the amyloidosis associated with carpal tunnel syndrome, senile

cardiac amyloidosis, familial amyloidotic polyneuropathy, and the amyloidosis associated with endocrine tumors, and especially is Alzheimer's disease.


In a third aspect, this invention provides a method of treating a disease characterized by  $\alpha$ -synuclein fibril formation in a mammal suffering therefrom, comprising administration to the mammal of a therapeutically effective amount of an isolated pure compound selected from the group consisting of the compounds of formula A, formula B, formula C, formula D, and formula E:




Formula A




Formula B



Formula C



Formula D



Formula E

where:

R is selected from the group consisting of hydrogen, 2,3-dihydroxybenzoyl, 3,4-dihydroxybenzoyl, 2,3,4-trihydroxybenzoyl, and 3,4,5-trihydroxybenzoyl;

R' is hydrogen or OH;

R<sub>1</sub> and R<sub>2</sub> are independently selected from hydrogen and non-interfering substituents;

X is selected from hydrogen and the group consisting of

- (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,
- (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
- (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents,
- (d) sugars, optionally substituted with one or more anionic groups selected from sulfate, phosphate, phosphonate, carboxylate, and sulfonate groups,
- (e) peptides and peptide derivatives, and
- (f) -C(O)R<sub>3</sub> and -C(O)OR<sub>3</sub> (where R<sub>3</sub> is selected from the group consisting of (a) through (e) above); and

Y is hydrogen, hydroxy, C<sub>1-6</sub> alkoxy, benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl), or —OSO<sub>2</sub>R<sub>4</sub> (where R<sub>4</sub> is C<sub>1-6</sub> alkyl or phenyl optionally substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl);

and the group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallol, anthralin, anthrarobin, anthrurufin, apigenin, apigetrin, apiose, baicalein, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose, centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, chrysamminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromycetin, collinomycin, curvularin, cyanidin, cyanidin 3-glucoside, cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside, cyanidin 3-sophoroside, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin, diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscine, fustin, galangin, gallein, gallocyanine, gardenin A, gardenin B, gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecycline, hematein,

hydroxysophorobioside, hydroxysophoricoside, icariin, isoquercitrin, kaempferol, kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D, leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin, oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin, purpurogallin, quercetagenin, quercimritrin, quinalizarin, quinic acid, resistomycin, rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B, robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone, tiron, troxerutin, and tunichrome B1, and the pharmaceutically acceptable salts thereof.

10 In preferred embodiments of this third aspect, only one such compound is administered; the mammal is a human; and the disease is Lewy body disease or Parkinson's disease, especially Parkinson's disease.

In a fourth aspect, this invention provides a drug product for the treatment of a disease characterized by  $\alpha$ -synuclein fibril formation in a mammal suffering therefrom, comprising a container labeled or accompanied by a label indicating that the drug product is for the treatment of a disease characterized by  $\alpha$ -synuclein fibril formation, the container containing one or more dosage units each comprising at least one pharmaceutically acceptable excipient and, as an active ingredient, an isolated pure compound selected from those used in the method of the third aspect of this invention.

20 In preferred embodiments of this fourth aspect, the drug product contains only one such compound, the mammal is a human; and the disease is Lewy body disease or Parkinson's disease, especially Parkinson's disease.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

25

##### Definitions

"Alkyl" means a linear hydrocarbyl group having from one to the number of carbon atoms specified, or a branched or cyclic hydrocarbyl group having from three to the number of carbon atoms specified. "Alkyl" in this application is given a broader meaning than is conventional in organic chemistry and includes both saturated groups (those conventionally known as alkyl groups), monounsaturated groups (such as those conventionally known as alkenyl and alkynyl groups), and polyunsaturated groups,

30

except that the terms does not include groups containing aromatic moieties, as the term "aromatic" is conventionally used. Exemplary C<sub>1-6</sub> alkyl groups include methyl, ethyl, isopropyl, cyclopropyl, *tert*-butyl, cyclopropylmethyl, and hexyl.

An "aromatic" group is a cyclic (monocyclic, condensed bicyclic, or linked  
5 bicyclic) group having from 5 to 12 ring carbon atoms, and sufficient ring unsaturation that the group is "aromatic" as that term is conventionally used. Exemplary aromatic groups include phenyl, naphthyl, and biphenyl. A "heteroaromatic" group is an "aromatic" group as just defined in which from 1 to 4 of the ring carbon atoms have been  
10 replaced by O, S, or NR (where R is hydrogen or C<sub>1-6</sub> alkyl). Exemplary heteroaromatic groups include pyrrolyl, furanyl, thiophenyl, benzofuranyl, indolyl, and the like. Such aromatic and heteroaromatic groups may optionally be substituted with 1 or more, especially 1 to 3, non-interfering substituents.

An "isolated pure compound" is a compound in isolated purified form such as is  
15 conventional for active ingredients in the pharmaceutical industry, and specifically excludes the compound when found as a component in a mixture such as within a plant or part thereof, or an extract or decoction of such plant or part, even when such mixtures are partially purified to limit the number of components present therein. However, treatment with an "isolated pure compound" is not limited to treatment with the compound alone but also includes treatment with the compound when present in a pharmaceutical  
20 composition of the type conventional in pharmaceutical practice, i.e. including one or more pharmaceutical excipients; however it specifically excludes treatment with the compound when the compound is found as a component in a mixture such as within a plant or part thereof, or an extract or decoction of such plant or part, even when such mixtures are partially purified to limit the number of components present therein.

25 "Mammal" includes humans and non-human mammals, such as companion animals (cats, dogs, and the like) and farm animals (cattle, horses, sheep, goats, swine, and the like).

A "non-interfering substituent" is a substituent that, when present in a compound, does not adversely affect the pharmacological activity of the compound and is not  
30 pharmaceutically undesirable. Suitable non-interfering substituents include halogen and C<sub>1-6</sub> alkyl and C<sub>1-6</sub> alkoxy, each optionally substituted with up to 5 halogen atoms.

“Pharmaceutically acceptable excipient” means an excipient that is useful in preparing a pharmaceutical composition that is generally safe, non-toxic, and desirable, and includes excipients that are acceptable for veterinary use as well as for human pharmaceutical use. Such excipients may be solid, liquid, semisolid, or, in the case of an aerosol composition, gaseous.

“Pharmaceutically acceptable salts” means salts that are pharmaceutically acceptable and have the desired pharmacological properties. Such salts include salts that may be formed where acidic protons present in the compounds are capable of reacting with inorganic or organic bases. Suitable inorganic salts include those formed with the alkali metals, e.g. sodium and potassium, magnesium, calcium, and aluminum. Suitable organic salts include those formed with organic bases such as the amine bases, e.g. ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine, and the like. Such salts also include acid addition salts formed with inorganic acids (e.g. hydrochloric and hydrobromic acids) and organic acids (e.g. acetic acid, citric acid, maleic acid, and the alkane- and arene-sulfonic acids such as methanesulfonic acid and benzenesulfonic acid). When there are two acidic groups present, a pharmaceutically acceptable salt may be a mono-acid-mono-salt or a di-salt; and similarly where there are more than two acidic groups present, some or all of such groups can be salified.

A “protecting group” has the meaning conventionally associated with it in organic synthesis, i.e. a group that selectively blocks one or more reactive sites in a multifunctional compound such that a chemical reaction can be carried out selectively on another unprotected reactive site and such that the group can readily be removed after the selective reaction is complete.

A “therapeutically effective amount” means the amount that, when administered to an animal for treating a disease, is sufficient to effect treatment for the disease.

“Treating” or “treatment” of the disease includes preventing the disease from occurring in a mammal that may be predisposed to the disease but does not yet experience or exhibit symptoms of the disease (prophylactic treatment), inhibiting the disease (slowing or arresting its development), providing relief from the symptoms or side-effects of the disease (including palliative treatment), and relieving the disease (causing regression of the disease). “Treating” amyloidosis includes any one or more of



the following: preventing, inhibiting, reducing, disassembling, disrupting, and disaggregating amyloid fibrils and amyloid protein deposits, such as A $\beta$  and the other amyloids referred to in the BACKGROUND TO THE INVENTION. "Treating" an  $\alpha$ -synuclein fibril disease includes any one or more of the following: preventing, inhibiting, reducing, disassembling, disrupting, and disaggregating  $\alpha$ -synuclein fibrils and  $\alpha$ -synuclein-associated protein deposits, such as those in Lewy body disease and Parkinson's disease.

The compounds found in the compositions and used in the methods of this invention may possess one or more chiral centers, and can therefore be produced as individual stereoisomers or as mixtures of stereoisomers, depending on whether individual stereoisomers or mixtures of stereoisomers of the starting materials are used. Unless indicated otherwise, the description or naming of a compound or group of compounds is intended to include both the individual stereoisomers or mixtures (racemic or otherwise) of stereoisomers. Methods for the determination of stereochemistry and the separation of stereoisomers are well known to a person of ordinary skill in the art [see the discussion in Chapter 4 of March J: Advanced Organic Chemistry, 4th ed. John Wiley and Sons, New York, NY, 1992].

#### Presently Preferred Compounds

While the broadest definition of the invention is set out in the Summary of the Invention, certain compounds of this invention are presently preferred.

Presently preferred compounds of this invention are compounds where:

- R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of hydrogen; C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, and C<sub>1-6</sub> alkylthio (in each of which the alkyl group is optionally substituted with 1 to 5 halogen atoms); and halo;
- X is selected from hydrogen and the group consisting of
- (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,
  - (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,

(c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents, and  
(d)  $-C(O)R_3$  and  $-C(O)OR_3$  (where  $R_3$  is selected from the group consisting of (a) through (c) above),

5 especially where X is selected from hydrogen and the group consisting of hydroxy, amino,  $-C(O)R_3$ , and  $-C(O)OR_3$  (where  $R_3$  is selected from hydroxy, amino,  $C_{1-6}$  alkyl optionally substituted with 1 to 5 halogen atoms, and aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups and optionally substituted with 1 to 5 non-interfering substituents selected from  
10 halogen atoms and  $C_{1-6}$  alkyl and  $C_{1-6}$  alkoxy, each optionally substituted with 1 to 5 halogen atoms; and

Y is selected from the group consisting of hydrogen, hydroxy,  $C_{1-6}$  alkoxy, and benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl and  $C_{1-6}$  alkoxy, each optionally substituted with 1 to 5  
15 halogen atoms).

and their individual stereoisomers, and the pharmaceutically acceptable salts thereof.

Preferred compounds include the compounds of formula A and formula B, the compounds of formula C, the compounds of formula D, the compounds of formula E, and the compounds of the list given following the descriptions of the formulae in the first  
20 aspect of the invention within the BRIEF SUMMARY OF THE INVENTION.

A number of different preferences have been given above, and following any one of these preferences results in a compound or the composition or method of this invention that is more presently preferred than a compound in which that particular preference is not followed. However, these preferences are generally independent and additive; and  
25 following more than one of these preferences may result in a more presently preferred compound than one in which fewer of the preferences are followed.

Presently preferred compounds of this invention include 1,2,4-benzenetriol, ellagic acid, ethyl gallate, exifone, gallamide, gallic acid, 5-hydroxydopamine, myricetin, phloroglucide, propyl gallate, quercetin, quinic acid, and tannic acid.

**Pharmacology and Utility**

The compounds of this invention act to inhibit or prevent amyloid fibril formation, inhibit or prevent amyloid fibril growth, and/or cause disassembly, disruption, and/or disaggregation of preformed amyloid fibrils and amyloid protein deposits. Their activity can be measured *in vitro* by methods such as those discussed in Examples 1 through 4 and Assay 1 below, while their activity *in vivo* against amyloidoses can be measured in animal models, such as those of Alzheimer's disease and in humans by a method such as that discussed in Assay 2 below.

The compounds of this invention also act to inhibit or prevent  $\alpha$ -synuclein fibril formation, inhibit or prevent  $\alpha$ -synuclein fibril growth, and/or cause disassembly, disruption, and/or disaggregation of preformed  $\alpha$ -synuclein fibrils and  $\alpha$ -synuclein-associated protein deposits. Their activity can be measured *in vitro* by methods similar to those discussed in Examples 1 through 4 below.

The therapeutic ratio of a compound can be determined, for example, by comparing the dose that gives effective anti-fibril (anti-amyloid or anti- $\alpha$ -synuclein) activity in a suitable *in vivo* model in a suitable animal species such as the mouse, with the dose that gives significant weight loss (or other observable side-effects) in the test animal species.

**Pharmaceutical compositions and administration**

In general, compounds of this invention will be administered in pure isolated form in therapeutically effective amounts by any of the usual modes known in the art, either singly or in combination with at least one other compound of this invention and/or at least one other conventional therapeutic agent for the disease being treated. A therapeutically effective amount may vary widely depending on the disease, its severity, the age and relative health of the animal being treated, the potency of the compound(s), and other factors. As anti-fibril agents, therapeutically effective amounts of compounds of this invention may range from 1-1000 mg/Kg body weight; for example, 10-100 mg/Kg. A person of ordinary skill in the art will be able without undue experimentation, having regard to that skill and this disclosure, to determine a therapeutically effective amount of a compound of this invention for the treatment of amyloidosis.

In general, compounds of this invention will be administered as pharmaceutical compositions by one of the following routes: oral, topical, systemic (e.g. transdermal, intranasal, or by suppository), or parenteral (e.g. intramuscular, subcutaneous, or intravenous injection). Compositions may take the form of tablets, pills, capsules, semisolids, powders, sustained release formulations, solutions, suspensions, elixirs, aerosols, or any other appropriate compositions; and comprise at least one compound of this invention in combination with at least one pharmaceutically acceptable excipient. Suitable excipients are well known to persons of ordinary skill in the art, and they, and the methods of formulating the compositions, may be found in such standard references as Alfonso AR: Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton PA, 1985. Suitable liquid carriers, especially for injectable solutions, include water, aqueous saline solution, aqueous dextrose solution, and glycols.

In particular, the compound(s) — preferably only one such compound is administered in any particular dosage form — can be administered, orally, for example, as tablets, troches, lozenges, aqueous or oily suspension, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known in the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations.

Tablets contain the compound in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, maize starch or alginic acid; binding agents, for example, maize starch, gelatin or acacia, and lubricating agents, for example, magnesium stearate or stearic acid or tale. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glycerol monostearate or glycerol distearate may be employed. Formulations for oral use may also

be presented as hard gelatin capsules wherein the compound is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin or olive oil.

- 5           Aqueous suspensions contain the compound in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethyl cellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be naturally occurring phosphatides, for example
- 10   lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example, heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids such as hexitol such as
- 15   polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters from fatty acids and a hexitol anhydrides, for example, polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more
- 20   preservatives, for example, ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, or one or more sweetening agents, such as sucrose or saccharin.
- 20           Oily suspensions may be formulated by suspending the compound in a vegetable oil, for example arachis oil, olive oil, sesame oil, or coconut oil or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example
- 25   beeswax, hard paraffin or cetyl alcohol. Sweetening agents, such as those set forth below, and flavoring agents may be added to provide a palatable oral preparation. These
- 25   compositions may be preserved by the addition of an antioxidant such as ascorbic acid. Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or
- 30   wetting agent, a suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already described above.
- 30   Additional excipients, for example sweetening, flavoring and agents, may also be present.

The compounds may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oils, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally occurring phosphatides, for example soy bean, lecithin, and occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsion may also contain sweetening and flavoring agents. Syrups and elixirs may be formulated with sweetening agents, for example, glycerol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

The compound can also be administered by injection or infusion, either subcutaneously or intravenously, or intramuscularly, or intrasternally, or intranasally, or by infusion techniques in the form of sterile injectable or oleaginous suspension. The compound may be in the form of a sterile injectable aqueous or oleaginous suspensions. These suspensions may be formulated according to the known art using suitable dispersing of wetting agents and suspending agents which have been described above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oils may be conventionally employed including synthetic mono- or diglycerides. In addition fatty acids such as oleic acid find use in the preparation of injectables.

Dosage regimens can be adjusted to provide the optimum therapeutic response. For example, several divided dosages may be administered daily or the dosage may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

It is especially advantageous to formulate the compounds in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each

containing a therapeutically effective quantity of the compound and at least one pharmaceutical excipient. A drug product will comprise a dosage unit form within a container that is labeled or accompanied by a label indicating the intended method of treatment, such as the treatment of an amyloid disease, such as Alzheimer's disease, or of a disease associated with  $\alpha$ -synuclein fibril formation, such as Parkinson's disease. A "therapeutically effective dosage" preferably inhibits amyloidosis or a disease associated with  $\alpha$ -synuclein fibril formation in a patient by at least 20, more preferably by at least 40%, even more preferably by at least 60%, and still more preferably by at least 80%, relative to untreated subjects.

10

#### **Preparation of the Compounds of this Invention**

Many of the compounds used in the compositions and methods of this invention are well known to the art. They may be briefly described in such references as the Merck Index, 12th edition, Merck & Co., Inc., Whitehouse Station, New Jersey, 1996 (which typically provides a reference to a synthesis or isolation), and may be found in chemical catalogs, such as those of commercial suppliers such as Aldrich Chemical Company (Milwaukee, WI), Bachem (Torrance, CA), Sigma (St. Louis, MO).

For those compounds that are novel, the starting materials and reagents used in preparing these compounds are generally available from commercial suppliers such as Aldrich Chemical Company, Bachem, and Sigma, or are prepared by methods well known to a person of ordinary skill in the art following procedures described in such references as Fieser and Fieser's Reagents for Organic Synthesis, vols 1-17, John Wiley and Sons, New York, NY, 1991; Rodd's Chemistry of Carbon Compounds, vols. 1-5 and supps, Elsevier Science Publishers, 1989; Organic Reactions, vols 1-40, John Wiley and Sons, New York, NY, 1991; March J: Advanced Organic Chemistry, 4th ed. John Wiley and Sons, New York, NY, 1992; and Larock: Comprehensive Organic Transformations, VCH Publishers, 1989, and the syntheses of the novel compounds will be readily suggested to a person of ordinary skill in the art by reference to known analogs (such as the commercially available analogs referred to above) of the novel compounds. Many such preparations will involve the use of protecting groups, especially for the protection of the hydroxy groups that form an essential part of the compounds; and the knowledge

and use of such protecting groups will be within the knowledge of a person of ordinary skill in the art.

The starting materials, intermediates, and compounds of this invention may be isolated and purified using conventional techniques, including filtration, distillation, crystallization, chromatography, and the like. They may be characterized using conventional methods, including physical constants and spectral data.

#### Examples

The following non-limiting examples illustrate the invention.

#### 10 **Example 1. Disassembly/disruption of Alzheimer's disease A $\beta$ 1-42 fibrils by polyhydroxylated aromatic compounds**

In this study, different types of commercially available compounds which consist of various polyhydroxylated aromatic containing structures were tested for their ability to cause a disassembly/disruption of pre-formed Alzheimer's disease amyloid fibrils containing A $\beta$  1-42. This type of activity would be important for any potential anti-amyloid drug which can be used in patients who already have substantial amyloid deposition in organs and/or tissues. For example, Alzheimer's disease patients in mid-to-late stage disease have abundant A $\beta$ -containing amyloid deposits in their brains as part of both neuritic plaques and cerebrovascular amyloid deposits. A compound capable of causing disassembly/disruption of pre-existing amyloid deposits would be advantageous for use in these patients who are at latter stages of the disease process.

For the first study, 1 mg of A $\beta$  1-42 (Bachem Inc., Torrance, CA, USA) was dissolved in 1.0 ml of double distilled water (1 mg/ml solution). 25  $\mu$ M of A $\beta$  1-42 was then incubated overnight (~18 hours) at 37°C, in the absence or presence of 100 $\mu$ g/ml of the following compounds: 1) EDTA (Sigma Chemical Company, St. Louis, MO, USA), 2) myricetin (Acros, Somerville, New Jersey, USA), 3) exifone (Acros) 4) pyrogallol (Sigma), 5) tannic acid (Acros), 6) pyrocatechol (Acros), 7) quercetin (Sigma), 8) ellagic acid (Acros), 9) 1,2,4-benzenetriol (Acros), 10) 5-hydroxydopamine (Acros), 11) gallamide hydrate (Acros), 12) gallic acid (Sigma), 13) ethyl gallate (Acros), 14) quinic acid (Acros), 15) propyl gallate (Sigma), and 16) phloroglucide (Acros), each in the



presence of 150 mM Tris HCl, 10 mM NaCl (pH 7.0) with 0.02% sodium azide. In this study, the A $\beta$  1-42:compound weight ratio was 1:1.

For the second study, 1 mg of A $\beta$  1-42 (Bachem) was dissolved in 1.0 ml of double distilled water (1 mg/ml solution). 25  $\mu$ M of A $\beta$  1-42 was then incubated  
5 overnight (~18 hours) at 37°C, in the absence or presence of 50 $\mu$ g/ml of the following compounds: 1) gallic acid, 2) ethyl gallate, 3) quinic acid, 4) gallamide trihydrate, 5) ellagic acid, 6) propyl gallate, and 7) pyrogallol, each in the presence of 150 mM Tris HCl, 10 mM NaCl (pH 7.0) with 0.02% sodium azide. In this study, the A $\beta$  1-42:compound weight ratio was 2:1.

10 A previously described method of measuring amyloid fibril formation utilizing Thioflavin T fluorometry (H Naiki et al., *Lab. Invest.* 65:104-110, 1991; H Levine III, *Protein Sci.* 2:404-410, 1993; H Levine III, *Amyloid: Int. J. Exp. Clin. Invest.* 2:1-6, 1995; H Naiki and K. Nakakuki, *Lab. Invest.* 74:374-383, 1996) was employed to identify potential therapeutic compounds capable of causing a disassembly/disruption of  
15 Alzheimer's A $\beta$  1-42 amyloid fibrils. Thioflavin T is known to bind to fibrillar amyloid proteins, and an increase in fluorescence correlates with an increase in amyloid fibril formation, whereas a decrease in fluorescence correlates with a decrease in amyloid fibrils due to disassembly and/or disruption. The Alzheimer's A $\beta$  protein (1-42) when placed in solution, such as distilled water, tends to spontaneously form amyloid fibrils.  
20 Using this sensitive assay, any decreases or increases in fluorescence was previously shown to correlate with a decrease or increase in the amount of amyloid fibrils (see the documents cited above), allowing one to identify and quantitate the extent of potential inhibitors and/or enhancers of Alzheimer's A $\beta$  1-42 amyloid fibrils.

To assess the effects of each compound on potential disassembly/ disruption of  
25 preformed A $\beta$  1-42 fibrils, 50  $\mu$ l of A $\beta$  1-42 with or without test compounds (described above) were added to 1.2 ml of 100 $\mu$ M Thioflavin T (Sigma) in 50mM NaH<sub>2</sub>PO<sub>4</sub> (pH 6.0) for fluorometry readings. Studies indicated that increasing concentrations of A $\beta$  gave a proportional increase in fluorescence in the presence of 100 $\mu$ M Thioflavin T, ruling out the presence of any disproportionate inner filter effects in these studies. Fluorescence  
30 emission at 482 nm was measured on a Turner instrument-model 450 fluorometer at an excitation wavelength of 450 nm. For each determination, the fluorometer was calibrated

by zeroing in the presence of the Thioflavin T reagent alone, and by setting the 50 ng/ml riboflavin (Sigma Chemical Co., St. Louis, MO) in the Thioflavin T reagent to 1800 fluorescence units. All fluorescence determinations were based on these references and any fluorescence given off by any of the compounds in the presence of the Thioflavin T reagent was always subtracted from all pertinent readings.

For all fibrillogenesis studies utilizing Thioflavin T fluorometry, as disclosed herein, comparisons of amyloid protein in the presence or absence of test compounds were based on paired Student's *t* tests with data shown as the mean of triplicate measurements  $\pm$  standard deviation.

As shown in Table 1, the polyhydroxylated aromatic compounds caused a disassembly/disruption of A $\beta$  1-42 amyloid fibril as determined by inhibition of Thioflavin T fluorescence. All results were significant at the  $p < 0.005$  level, except that for quinic acid at the 2:1 ratio (asterisked in Table 1), which was not significant.

Table 1: Disassembly/disruption of Alzheimer's 1-42 fibrils, as indicated by Thioflavin T fluorescence inhibition

Compound name	Fluorescence inhibition, %, at the A $\beta$ 1-42:compound w/w ratios given	
	1:1	2:1
Myricetin	94 $\pm$ 0.9	
Exifone	93 $\pm$ 1.4	
Pyrogallol	89 $\pm$ 6.7	72 $\pm$ 3.8
Tannic acid	77 $\pm$ 1.3	
Pyrocatechol	77 $\pm$ 2.6	
Quercetin	76 $\pm$ 0.6	
Ellagic acid	74 $\pm$ 1.4	62 $\pm$ 3.9
1,2,4-Benzenetriol	71 $\pm$ 3.3	
5-Hydroxydopamine	70 $\pm$ 1.1	
Gallamide trihydrate	65 $\pm$ 12.3	60 $\pm$ 2.2
Gallic acid	57 $\pm$ 1.9	44 $\pm$ 1.7
Ethyl gallate	49 $\pm$ 0.8	30 $\pm$ 3.7

Quinic acid	31 ± 9.0	0.5 ± 3.9*
Phloroglucide	30 ± 0.6	
Propyl gallate	29 ± 2.8	38 ± 4.8

EDTA, a known chelating agent, caused no significant disassembly/disruption of A $\beta$  1-42 amyloid fibrils, suggesting that the inhibitory effects observed with polyhydroxylated aromatic compounds was not attributable to their ability to complex metals.

**Example 2. Dose-dependent disassembly/disruption of Alzheimer's disease A $\beta$  1-40 fibrils by tannic acid and gallic acid**

In this study, the potential dose-dependent effects of tannic acid and gallic acid on disassembly/ disruption of pre-formed A $\beta$  1-40 was assessed. In this experiment, 1 mg of A $\beta$  1-40 (Bachem Inc., Torrance, CA, USA; Lot # T-20824) was dissolved in 1.0 ml of double distilled water (1 mg/ml solution) and incubated for 4 days at 37°C to spontaneously induce fibril formation. 25  $\mu$ M of pre-fibrillized A $\beta$  1-40 was then incubated overnight (~18 hours) at 37°C, in the absence or presence of increasing amounts (25 $\mu$ g/ml, 50 $\mu$ g/ml, 75 $\mu$ g/ml and 100 $\mu$ g/ml) of tannic acid or gallic acid (each in the presence of 150 mM Tris HCl, 10 mM NaCl, pH 7.0, with 0.02% sodium azide). The A $\beta$ :compound weight ratios were therefore 4:1, 2:1, 4:3, and 1:1, respectively. 50  $\mu$ l aliquots were then added to 1.2 ml of 100 $\mu$ M Thioflavin T (Sigma) in 50mM NaH<sub>2</sub>PO<sub>4</sub> (pH 6.0) for fluorometry readings as described in Example 1 above.

As shown in Table 2, both tannic acid and gallic acid caused a dose-dependent disassembly/ disruption of A $\beta$  1-40 amyloid fibrils as indicated by a dose-dependent inhibition of Thioflavin T fluorescence. All results were significant at the  $p < 0.005$  level, except that for gallic acid at the 4:1 ratio (asterisked in Table 2), which was significant at the  $p < 0.05$  level.

Table 2: Dose-dependent disassembly/disruption of Alzheimer's 1-40 fibrils, as indicated by Thioflavin T fluorescence inhibition

Compound name	Fluorescence inhibition, %, at the A $\beta$ 1-40:compound w/w ratios given			
	4:1	2:1	4:3	1:1
Tannic acid	31 $\pm$ 4.8	42 $\pm$ 2.8	49 $\pm$ 3.7	53 $\pm$ 4.2
Gallic acid	14 $\pm$ 8.2*	22 $\pm$ 3.3	34 $\pm$ 3.6	45 $\pm$ 4.1

**5 Example 3. Disaggregation of Alzheimer's disease A $\beta$  1-40 fibrils by polyhydroxylated aromatic compounds**

In this study, a Congo red-A $\beta$  spectrophotometric assay (Klunk et al., *Anal. Biochem.* 266:66-76, 1999) was modified to determine the effectiveness of polyhydroxylated aromatic compounds on the disaggregation of pre-formed A $\beta$  1-40 amyloid fibrils. For this assay, 1 mg of A $\beta$  1-40 (Bachem) was incubated for 4 days in distilled water at 37°C to spontaneously produce amyloid fibrils. 25  $\mu$ M of fibrillized A $\beta$  1-40 was then incubated in triplicate with various test compounds for 3 days at 37°C in Tris-buffered saline (TBS) (100 mM Tris; 50 mM NaCl; pH 7.0, with 0.02% sodium azide), at an A $\beta$ :compound weight ratio of 2:1. Following incubation, 50  $\mu$ l of 360  $\mu$ M Congo red (Sigma) in distilled water was then added to 250  $\mu$ l of each incubation mixture, giving a final A $\beta$ :Congo red molar ratio of 1:3. After 10 minutes, the absorbance at 405 nm (reference wavelength to account for the absorbance of Congo red alone at 540 nm) and 540 nm (sample absorbance where "sample" refers to A $\beta$  alone, test compound alone, or A $\beta$  plus test compound, all in the presence of Congo red) was determined using a Biorad Model 550 ELISA Plate Reader (Biorad, Hercules, CA, USA). The absorbance at wavelength 405 nm was automatically subtracted by the ELISA plate reader from the absorbance at wavelength 540 nm (difference is referred to as  $\Delta$  absorbance) (see Klunk et al. cited above). Therefore, the  $\Delta$  absorbance reading at 540 nm was proportional to the amount of aggregated A $\beta$  left in solution (Klunk et al.).

25 For all experiments involving test compounds, the  $\Delta$  absorbance reading at 540 nm of the test compound alone (in the absence of A $\beta$ ), was always subtracted from the corresponding  $\Delta$  absorbance reading at 540 nm of the test compound in the presence of

A $\beta$ . Using this modification of the method of Klunk et al., the use of a greater final concentration of Congo red, i.e. 60 $\mu$ M instead of 14 $\mu$ M, in the presence of fibrillar A $\beta$  gave an overall absorbance at 540 nm that was always below 1.0 Absorbance Unit (AU), and well within the linear absorbance range.

5        The following polyhydroxylated aromatic containing compounds were tested using the above described Congo red-A $\beta$  spectrophotometric assay to determine their effectiveness on disaggregation of pre-formed A $\beta$  1-40 amyloid fibrils: 1) gallic acid, 2) ethyl gallate, 3) quinic acid, 4) gallamide trihydrate, 5) ellagic acid, 6) propyl gallate, and 7) pyrogallol.

10        The polyhydroxylated aromatic compounds had varying effects on causing disaggregation of pre-aggregated A $\beta$  1-40 amyloid fibrils as determined using the Congo red spectrophotometric assay described above. The results were significant at the  $p < 0.005$  level, except for propyl gallate (asterisked in Table 3) at the  $p < 0.05$  level, and quinic acid (double asterisked), which was not significant.

15        Table 3: Disaggregation of Alzheimer's 1-40 fibrils,  
as indicated by Congo red spectrophotometry

Compound name	Decrease in absorbance, %
Gallic acid	52 $\pm$ 0.4
Ethyl gallate	28 $\pm$ 5.0
Quinic acid	0 + 5.0**
Gallamide trihydrate	31 $\pm$ 1.9
Ellagic acid	54 $\pm$ 2.8
Propyl gallate	17 $\pm$ 7.3*
Pyrogallol	63 $\pm$ 3.4

20        **Example 4. Dose-dependent disaggregation of Alzheimer's disease A $\beta$  1-40 fibrils by tannic acid and gallic acid**

In this study, the potential dose-dependent effects of tannic acid and gallic acid on the disaggregation of fibrillized A $\beta$  1-40 was assessed. In this experiment, the modified

Congo red-A $\beta$  spectrophotometric assay (Klunk et al, *Anal. Biochem.* 266:66-76, 1999) was used as described above (i.e. Example 4). However, in this specific experiment increasing amounts of tannic acid or gallic acid (i.e. 25 $\mu$ g/ml, 50 $\mu$ g/ml, 75 $\mu$ g/ml and 100 $\mu$ g/ml) were tested following an overnight (~18 hours) incubation at 37°C in the presence of 25 $\mu$ M of A $\beta$  1-40 (Bachem).

As shown in Table 4, both tannic acid and gallic acid caused a dose-dependent disaggregation of A $\beta$  1-40 amyloid fibril as determined by decreases in Thioflavin T fluorescence. All results were significant at the  $p < 0.001$  level, except that for gallic acid at the 4:1 ratio (asterisked in Table 4), which was significant at the  $p < 0.05$  level.

Table 4: Dose-dependent disaggregation of Alzheimer's 1-40 fibrils, as indicated by Congo red spectrophotometry

Compound name	Decrease in absorbance, %, at the A $\beta$ 1-42: compound w/w ratios given			
	4:1	2:1	4:3	1:1
Tannic acid	42 $\pm$ 5.2	48 $\pm$ 6.8	59 $\pm$ 6.4	61 $\pm$ 11.1
Gallic acid	17 $\pm$ 9.5*	22 $\pm$ 4.0	30 $\pm$ 6.0	32 $\pm$ 4.8

#### Example 5. Disassembly/disruption of islet amyloid fibrils (amylin) by polyhydroxylated aromatic compounds

90% of patients with type II diabetes demonstrate the deposition and accumulation of amyloid fibrils in the islets of Langerhans in the pancreas (Cooper et al., *Proc. Natl. Acad. Sci. USA* 84:8628-8632, 1987). This amyloid protein involved consists of a 37 amino acid protein known as islet amyloid polypeptide or amylin. Islet amyloid is believed to contribute to the destruction of the beta-cells of the pancreas, thus eventually leading many patients to become insulin-dependent (i.e. type I diabetes). Amylin has the ability to also form substantial amyloid fibrils immediately when placed in solution. The next study was therefore implemented to determine whether some of the specific polyhydroxylated aromatic containing compounds which cause a disassembly/disruption of A $\beta$  fibrils, also cause a disassembly/disruption of islet amyloid fibrils.

For this study, the method of Thioflavin T fluorometry as described in Example 1 was used. Briefly, 25  $\mu$ M of human amylin (Bachem) was incubated overnight (~18 hours) at 37°C, alone or in the presence of 100 $\mu$ g/ml of the following compounds: 1) exifone, 2) myricetin, and 3) tannic acid, each in the presence of 150 mM Tris HCl, 10 mM NaCl, pH 7.0, with 0.02% sodium azide, at an amylin:compound weight ratio of 1:1.

Following Thioflavin T fluorometry readings as described in Example 1, 5 $\mu$ l aliquots of amylin only, amylin + myricetin, amylin + exifone, and amylin + tannic acid were also taken, allowed to air dry overnight on gelatin-coated slides, and stained with Congo red as previously described (Castillo et al., *Diabetes* 47:612-620, 1998).

As shown in Table 5, the polyhydroxylated aromatic compounds which were very effective in causing a disassembly/disruption of A $\beta$  1-42 amyloid fibrils were also effective in causing a disassembly/disruption of islet amyloid fibrils. All results were significant at the  $p < 0.005$  level.

Table 5: Disassembly/disruption of amylin fibrils,  
as indicated by Thioflavin T fluorescence inhibition

Compound name	Fluorescence inhibition, %
Myricetin	97.4 $\pm$ 0.3
Exifone	99.1 $\pm$ 0.5
Tannic acid	83.8 $\pm$ 1.4

Congo red staining experiments confirmed the disassembly/disruption of amylin fibrils by polyhydroxylated aromatic compounds initially demonstrated by Thioflavin T fluorometry studies as described above. Congo red staining of amylin alone demonstrated positive staining (i.e. classic red/green birefringence as viewed under polarized light and indicative of amyloid) (Püchtler et al., *J. Histochem. Cytochem.* 10:355-364, 1962). In comparison, an overnight incubation with exifone, myricetin or tannic acid resulted in a marked decrease in Congo red staining, suggestive of an amylin fibril disassembly/disruption.

Further *in vitro* and *in vivo* assays may be used to test the compounds for their effectiveness in the treatment of Alzheimer's disease, such as those described in European Published Patent Application No. 0 659 418.

Stock solutions of peptides (1 mM) are freshly prepared in pyrogen-free sterile water and diluted to the indicated concentrations in defined culture media. Rat hippocampal cultures (10-14 days *in vitro*) are treated with peptides or vehicle for four days. The viability of the rat cortical cultures is visually assessed by phase contrast microscopy and quantified by measuring lactate dehydrogenase (LDH) released into the culture media.

10

#### Assay 1

Primary rat hippocampal neurons are cultured *in vitro* with standard cell culture techniques. Amyloid-beta (A $\beta$ ) peptide is added to cultured cells at a normally toxic concentration of 25-50  $\mu$ M. After 4 days of treatment, viability is assessed by measurement of lactate dehydrogenase (LDH) released into culture medium. Lactate dehydrogenase (LDH) is measured in 20  $\mu$ l aliquots of conditioned defined DMEM using a standard 340 nm kinetic LDH assay (Sigma Catalog Number #228-20) in a 96 well format. Assays are performed at 37°C in a PC-driven EL340 Microplate Biokinetics plate reader (Bio-Tek Instruments) using Delta Soft II software (v. 3.30B, BioMetallics, Inc.) for data analysis. Quality control standards containing normal and elevated levels of serum LDH (for example, Sigma Enzyme Controls 2N and 2E) are run with every assay. Results are expressed as units of LDH/L where 1 unit is defined as the amount of enzyme that will catalyze the formation of 1 micromole of nicotinamide adenine dinucleotide per minute under conditions of the assay. For protection studies, a compound of formula 1 is added to cultures prior to and/or concurrently with the amyloid- beta treatment.

25

Activity of the compounds is illustrated by a decrease in LDH released into the media (a neurotoxic indicator), as compared to control.

#### Assay 2

30

Five to fifty women are selected for a clinical study. The women are post-menopausal, i.e., have ceased menstruating for between 6 and 12 months prior to the



study's initiation, have been diagnosed with early stage Alzheimer's disease (AD), are expected to have worsening symptoms of AD within the study period, but are in good general health otherwise. The study has a placebo control group, i.e., the women are divided into two groups, one of which receives the compound of this invention and the other receives a placebo. The patients are benchmarked as to memory, cognition, reasoning, and other symptoms associated with AD. Women in the test group receive a therapeutic dose of the compound per day by the oral route. They continue this therapy for 6-36 months. Accurate records are kept as to the benchmarked symptoms in both groups and at the end of the study these results are compared. The results are compared both between members of each group and also the results for each patient are compared to the symptoms reported by each patient before the study began. Activity of the compound is illustrated by an attenuation of the typical cognitive decline and/or behavioral disruptions associated with AD.

Utility of the compounds is evidenced by activity in at least one of the above assays.


While this invention has been described in conjunction with specific embodiments and examples, it will be apparent to a person of ordinary skill in the art, having regard to this disclosure, that equivalents of the specifically disclosed materials and techniques will also be applicable to this invention; and such equivalents are intended to be included within the following claims.

## WHAT IS CLAIMED IS:


5

1. A method of treating a mammal suffering from amyloidosis or a disease characterized by  $\alpha$ -synuclein fibril formation, comprising administration to the mammal of a therapeutically effective amount of an isolated pure compound selected from the group consisting of the compounds of formula A, formula B, formula C, formula D, and


10 formula E:




Formula A




Formula B



Formula C



Formula D



Formula E

where:

- 15 R is selected from the group consisting of hydrogen, 2,3-dihydroxybenzoyl, 3,4-dihydroxybenzoyl, 2,3,4-trihydroxybenzoyl, and 3,4,5-trihydroxybenzoyl;  
 R' is hydrogen or OH;  
 R<sub>1</sub> and R<sub>2</sub> are independently selected from hydrogen and non-interfering substituents;

X is selected from hydrogen and the group consisting of

- (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,
- (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of  
5 halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
- (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents,
- (d) sugars, optionally substituted with one or more anionic groups selected from  
10 sulfate, phosphate, phosphonate, carboxylate, and sulfonate groups,
- (e) peptides and peptide derivatives, and
- (f) -C(O)R<sub>3</sub> and -C(O)OR<sub>3</sub> (where R<sub>3</sub> is selected from the group consisting of (a) through (e) above); and

Y is hydrogen, hydroxy, C<sub>1-6</sub> alkoxy, benzyloxy (where the phenyl group is optionally  
15 substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl), or —OSO<sub>2</sub>R<sub>4</sub> (where R<sub>4</sub> is C<sub>1-6</sub> alkyl or phenyl optionally substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl);

and the group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallo, anthralin,  
20 anthrarobin, anthraquinone, apigenin, apigenin, apiose, baicalein, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose, centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, chrysamminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromycesin, collinomycesin, curvularin, cyanidin, cyanidin 3-glucoside,  
25 cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside, cyanidin 3-sophorose, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin, diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscin, fustin,  
30 galangin, gallein, galloycyanine, gardenin A, gardenin B, gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecycline, hematein,

hydroxysophorobioside, hydroxysophoricoside, icariin, isoquercitrin, kaempferol, kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D, leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin, oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin, purpurogallin, quercetagenin, quercimritrin, quinalizarin, quinic acid, resistomycin, rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B, robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone, tiron, troxerutin, and tunichrome B1,

but excluding pyrogallol,  
and the pharmaceutically acceptable salts thereof.

2. The method of Claim 1 where the amyloidosis is selected from the group of diseases consisting of Alzheimer's disease, Down's syndrome, hereditary cerebral hemorrhage with amyloidosis of the Dutch type, the amyloidosis of chronic inflammation, the amyloidosis of malignancy and familial Mediterranean fever, the amyloidosis of multiple myeloma and B-cell dyscrasias, the amyloidosis of type II diabetes, the amyloidosis of the prion diseases, Creutzfeldt-Jakob disease, Gerstmann-Straussler syndrome, kuru, scrapie, the amyloidosis associated with long-term hemodialysis, the amyloidosis associated with carpal tunnel syndrome, senile cardiac amyloidosis, familial amyloidotic polyneuropathy, and the amyloidosis associated with endocrine tumors.

3. The method of Claim 2 where the amyloidosis is Alzheimer's disease.

4. The drug method of Claim 1 where the  $\alpha$ -synuclein fibril formation is Lewy body disease or Parkinson's disease.

5. The method of Claim 1 where  $R_1$  and  $R_2$  are independently selected from the group consisting of hydrogen;  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, and  $C_{1-6}$  alkylthio (in each of which the alkyl group is optionally substituted with 1 to 5 halogen atoms); and halo.

6. The method of Claim 1 where X is selected from hydrogen and the group consisting of
- (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,
  - (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
  - (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents, and
  - (d) -C(O)R<sub>3</sub> and -C(O)OR<sub>3</sub> (where R<sub>3</sub> is selected from the group consisting of (a) through (c) above).
7. The method of Claim 1 where X is selected from hydrogen and the group consisting of hydroxy, amino, -C(O)R<sub>3</sub>, and -C(O)OR<sub>3</sub> (where R<sub>3</sub> is selected from hydroxy, amino, C<sub>1-6</sub> alkyl optionally substituted with 1 to 5 halogen atoms, and aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups and optionally substituted with 1 to 5 non-interfering substituents selected from halogen atoms and C<sub>1-6</sub> alkyl and C<sub>1-6</sub> alkoxy, each optionally substituted with 1 to 5 halogen atoms).
8. The method of Claim 1 where Y is selected from the group consisting of hydrogen, hydroxy, C<sub>1-6</sub> alkoxy, and benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and C<sub>1-6</sub> alkyl and C<sub>1-6</sub> alkoxy, each optionally substituted with 1 to 5 halogen atoms).
9. The method of Claim 1 where the compound is a compound of formula A or formula B, or a pharmaceutically acceptable salt thereof.
10. The method of Claim 9 where the compound is selected from the group consisting of dibromogallic acid, digallic acid, ethyl gallate, exifone, fisetin, gallacetophenone, gallamide, gallic acid, α-glucogallin, β-glucogallin, 5-hydroxydopamine, and propyl gallate, and the pharmaceutically acceptable salts thereof.

11. The method of Claim 1 where the compound is a compound of formula C or a pharmaceutically acceptable salt thereof.
12. The method of Claim 1 where the compound is a compound of formula D or a  
5 pharmaceutically acceptable salt thereof.
13. The method of Claim 12 where the compound is ellagic acid or a pharmaceutically acceptable salt thereof.
- 10 14. The method of Claim 1 where the compound is a compound of formula E or a pharmaceutically acceptable salt thereof.
- 15 15. The method of Claim 14 where the compound is selected from the group consisting of catechin, epicatechin, gallic acid, epigallocatechin, and their gallate esters, and the pharmaceutically acceptable salts thereof.
16. The method of Claim 1 where the active ingredient is selected from group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallol, anthralin, anthrarobin, anthrakinone, apigenin,  
20 apigetrin, apiose, baicalin, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose, centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, chrysaminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromyctin, collinomyctin, curvularin, cyanidin, cyanidin 3-glucoside, cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside,  
25 cyanidin 3-sophoroside, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin, diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscine, fustin, galangin, gallein, gallocyanine, gardenin A, gardenin B,  
30 gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecycline, hemeatin, hydroxysophorobioside, hydroxysophoricoside, icariin, isoquercitrin,


kaempferol, kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D, leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin, oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin, purpurogallin, pyrocatechol, quercetagenin, quercimritrin, quinalizarin, quinic acid, resistomycin, 5 rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B, robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone, tiron, troxerutin, and tunichrome B1, and the pharmaceutically acceptable salts thereof.

17. The method of Claim 1 where the compound is selected from 1,2,4-benzenetriol, 10 ellagic acid, ethyl gallate, exifone, gallamide, gallic acid, 5-hydroxydopamine, myricetin, phloroglucide, propyl gallate, quercetin, quinic acid, and tannic acid, and the pharmaceutically acceptable salts thereof.


18. The method of Claim 17 where the compound is selected from the group 15 consisting of myricetin and quercetin, and the pharmaceutically acceptable salts thereof.

19. Pharmaceutical composition adapted for treating a mammal suffering from amyloidosis or a disease characterized by  $\alpha$ -synuclein fibril formation, comprising a therapeutically effective amount of an isolated pure compound selected from the group 20 consisting of the compounds of formula A, formula B, formula C, formula D, and formula E:


39




Formula A




Formula B



Formula C



Formula D



Formula E

where:

- 5 R is selected from the group consisting of hydrogen, 2,3-dihydroxybenzoyl, 3,4-dihydroxybenzoyl, 2,3,4-trihydroxybenzoyl; and 3,4,5-trihydroxybenzoyl;  
R' is hydrogen or OH;
- R<sub>1</sub> and R<sub>2</sub> are independently selected from hydrogen and non-interfering substituents;
- X is selected from hydrogen and the group consisting of
  - 10 (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,
  - (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
  - 15 (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents,



(d) sugars, optionally substituted with one or more anionic groups selected from sulfate, phosphate, phosphonate, carboxylate, and sulfonate groups,

(e) peptides and peptide derivatives, and

(f)  $-C(O)R_3$  and  $-C(O)OR_3$  (where  $R_3$  is selected from the group consisting of (a)

5 through (e) above); and

Y is hydrogen, hydroxy,  $C_{1-6}$  alkoxy, benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl), or  $-OSO_2R_4$  (where  $R_4$  is  $C_{1-6}$  alkyl or phenyl optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl);

10 and the group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallol, anthralin, anthrarobin, anthrarufin, apigenin, apigetrin, apiose, baicalein, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose, centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, 15 chrysamminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromycetin, collinomycin, curvularin, cyanidin, cyanidin 3-glucoside, cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside, cyanidin 3-sophoroside, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin, diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome 20 A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscine, fustin, galangin, gallein, gallocyanine, gardenin A, gardenin B, gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecycline, hematein, hydroxysophorobioside, hydroxysophoricoside, icariin, isoquercitrin, kaempferol, 25 kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D, leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin, oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin, purpurogallin, quercetagenin, quercimritrin, quinalizarin, quinic acid, resistomycin, rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B, 30 robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone, tiron, troxerutin, and tunichrome B1,

but excluding pyrogallol,  
and the pharmaceutically acceptable salts thereof.

20. The composition of Claim 19 where the amyloidosis is selected from the group of  
5 diseases consisting of Alzheimer's disease, Down's syndrome, hereditary cerebral  
hemorrhage with amyloidosis of the Dutch type, the amyloidosis of chronic  
inflammation, the amyloidosis of malignancy and familial Mediterranean fever, the  
amyloidosis of multiple myeloma and B-cell dyscrasias, the amyloidosis of type II  
diabetes, the amyloidosis of the prion diseases, Creutzfeldt-Jakob disease, Gerstmann-  
10 Straussler syndrome, kuru, scrapie, the amyloidosis associated with long-term  
hemodialysis, the amyloidosis associated with carpal tunnel syndrome, senile cardiac  
amyloidosis, familial amyloidotic polyneuropathy, and the amyloidosis associated with  
endocrine tumors.
- 15 21. The composition of Claim 20 where the amyloidosis is Alzheimer's disease.
22. The composition of Claim 19 where the  $\alpha$ -synuclein fibril formation is Lewy  
body disease or Parkinson's disease.
- 20 23. The composition of Claim 19 where  $R_1$  and  $R_2$  are independently selected from  
the group consisting of hydrogen;  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, and  $C_{1-6}$  alkylthio (in each of  
which the alkyl group is optionally substituted with 1 to 5 halogen atoms); and halo.
24. The composition of Claim 19 where X is selected from hydrogen and the group  
25 consisting of  
(a) hydroxy, amino,  $C_{1-6}$  alkylamino, di( $C_{1-6}$  alkyl)amino, and cycloamino,  
(b)  $C_{1-22}$  alkyl,  $C_{1-22}$  alkoxy,  $C_{1-22}$  alkylthio, and  $C_{1-22}$  alkylcarboxyl, each optionally  
substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy,  
mercapto, amino, nitro,  $C_{1-6}$  alkoxy,  $C_{1-6}$  alkylthio, and  $C_{1-6}$  alkylcarboxyl,  
30 (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups,  
and optionally substituted with 1 to 5 non-interfering substituents, and

(d)  $-C(O)R_3$  and  $-C(O)OR_3$  (where  $R_3$  is selected from the group consisting of (a) through (c) above).

25. The composition of Claim 19 where X is selected from hydrogen and the group consisting of hydroxy, amino,  $-C(O)R_3$ , and  $-C(O)OR_3$  (where  $R_3$  is selected from hydroxy, amino,  $C_{1-6}$  alkyl optionally substituted with 1 to 5 halogen atoms, and aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups and optionally substituted with 1 to 5 non-interfering substituents selected from halogen atoms and  $C_{1-6}$  alkyl and  $C_{1-6}$  alkoxy, each optionally substituted with 1 to 5 halogen atoms.


10

26. The composition of Claim 19 where Y is selected from the group consisting of hydrogen, hydroxy,  $C_{1-6}$  alkoxy, and benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl and  $C_{1-6}$  alkoxy, each optionally substituted with 1 to 5 halogen atoms).


15

27. Use of an isolated pure compound selected from the group consisting of the compounds of formula A, formula B, formula C, formula D, and formula E:


43




Formula A




Formula B



Formula C



Formula D



Formula E

where:

- 5 R is selected from the group consisting of hydrogen, 2,3-dihydroxybenzoyl, 3,4-dihydroxybenzoyl, 2,3,4-trihydroxybenzoyl, and 3,4,5-trihydroxybenzoyl;  
 R' is hydrogen or OH;  
 R<sub>1</sub> and R<sub>2</sub> are independently selected from hydrogen and non-interfering substituents;  
 X is selected from hydrogen and the group consisting of
- 10 (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,  
 (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
- 15 (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents,

- (d) sugars, optionally substituted with one or more anionic groups selected from sulfate, phosphate, phosphonate, carboxylate, and sulfonate groups,  
(e) peptides and peptide derivatives, and  
(f)  $-C(O)R_3$  and  $-C(O)OR_3$  (where  $R_3$  is selected from the group consisting of (a) through (e) above); and

Y is hydrogen, hydroxy,  $C_{1-6}$  alkoxy, benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl), or  $-OSO_2R_4$  (where  $R_4$  is  $C_{1-6}$  alkyl or phenyl optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl);

- and the group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallol, anthralin, anthrarobin, antharufin, apigenin, apigetrin, apiose, baicaléin, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose, centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, chrysamminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromycetin, collinomycin, curvularin, cyanidin, cyanidin 3-glucoside, cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside, cyanidin 3-sophoroside, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin, diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscine, fustin, galangin, gallein, gallocyanine, gardenin A, gardenin B, gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecyclyne, hematein, hydroxysophorobioside, hydroxysophoricoside, icariin, isoquercitrin, kaempferol, kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D, leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin, oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin, purpurogallin, quercetagenin, quercimritrin, quinalizarin, quinic acid, resistomycin, rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B, robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone, tiron, troxerutin, and tunichrome B1,

but excluding pyrogallol,  
and the pharmaceutically acceptable salts thereof, for the manufacture of a  
pharmaceutical preparation adapted for treating a mammal suffering from amyloidosis or  
a disease characterized by  $\alpha$ -synuclein fibril formation.

5

28. The use of Claim 27 where the amyloidosis is selected from the group of diseases  
consisting of Alzheimer's disease, Down's syndrome, hereditary cerebral hemorrhage  
with amyloidosis of the Dutch type, the amyloidosis of chronic inflammation, the  
amyloidosis of malignancy and familial Mediterranean fever, the amyloidosis of multiple  
10 myeloma and B-cell dyscrasias, the amyloidosis of type II diabetes, the amyloidosis of  
the prion diseases, Creutzfeldt-Jakob disease, Gerstmann-Straussler syndrome, kuru,  
scrapie, the amyloidosis associated with long-term hemodialysis, the amyloidosis  
associated with carpal tunnel syndrome, senile cardiac amyloidosis, familial amyloidotic  
polyneuropathy, and the amyloidosis associated with endocrine tumors.

15

29. The use of Claim 28 where the amyloidosis is Alzheimer's disease.

30. The use of Claim 27 where the  $\alpha$ -synuclein fibril formation is Lewy body disease  
or Parkinson's disease.

20

31. The use of Claim 27 where  $R_1$  and  $R_2$  are independently selected from the group  
consisting of hydrogen;  $C_{1-6}$  alkyl,  $C_{1-6}$  alkoxy, and  $C_{1-6}$  alkylthio (in each of which the  
alkyl group is optionally substituted with 1 to 5 halogen atoms); and halo.

25 32. The use of Claim 27 where X is selected from hydrogen and the group consisting  
of

(a) hydroxy, amino,  $C_{1-6}$  alkylamino, di( $C_{1-6}$  alkyl)amino, and cycloamino,

(b)  $C_{1-22}$  alkyl,  $C_{1-22}$  alkoxy,  $C_{1-22}$  alkylthio, and  $C_{1-22}$  alkylcarboxyl, each optionally  
substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy,

30 mercapto, amino, nitro,  $C_{1-6}$  alkoxy,  $C_{1-6}$  alkylthio, and  $C_{1-6}$  alkylcarboxyl,

(c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups,

and optionally substituted with 1 to 5 non-interfering substituents, and


(d)  $-C(O)R_3$  and  $-C(O)OR_3$  (where  $R_3$  is selected from the group consisting of (a) through (c) above).

- 5 33. The use of Claim 27 where X is selected from hydrogen and the group consisting of hydroxy, amino,  $-C(O)R_3$ , and  $-C(O)OR_3$  (where  $R_3$  is selected from hydroxy, amino,  $C_{1-6}$  alkyl optionally substituted with 1 to 5 halogen atoms, and aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups and optionally substituted with 1 to 5 non-interfering substituents selected from halogen atoms and  $C_{1-6}$  alkyl and  $C_{1-6}$  alkoxy, each optionally substituted with 1 to 5 halogen atoms.


- 10 34. The use of Claim 27 where Y is selected from the group consisting of hydrogen, hydroxy,  $C_{1-6}$  alkoxy, and benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl and  $C_{1-6}$  alkoxy, each optionally substituted with 1 to 5 halogen atoms).

- 15 35. A drug product for the treatment of a mammal suffering from amyloidosis or a disease characterized by  $\alpha$ -synuclein fibril formation, comprising a container labeled or accompanied by a label indicating that the drug product is for the treatment of amyloidosis or a disease characterized by  $\alpha$ -synuclein fibril formation, the container containing one or more dosage units each comprising at least one pharmaceutically acceptable excipient and, as an active ingredient, an isolated pure compound selected from the group consisting of the compounds of formula A, formula B, formula C, formula D, and formula E:
- 20


47




Formula A




Formula B



Formula C



Formula D



Formula E

where:

- 5 R is selected from the group consisting of hydrogen, 2,3-dihydroxybenzoyl, 3,4-dihydroxybenzoyl, 2,3,4-trihydroxybenzoyl, and 3,4,5-trihydroxybenzoyl;  
R' is hydrogen or OH;
- R<sub>1</sub> and R<sub>2</sub> are independently selected from hydrogen and non-interfering substituents;
- X is selected from hydrogen and the group consisting of
  - 10 (a) hydroxy, amino, C<sub>1-6</sub> alkylamino, di(C<sub>1-6</sub> alkyl)amino, and cycloamino,
  - (b) C<sub>1-22</sub> alkyl, C<sub>1-22</sub> alkoxy, C<sub>1-22</sub> alkylthio, and C<sub>1-22</sub> alkylcarboxyl, each optionally substituted with 1 to 5 moieties selected from the group consisting of halogen, hydroxy, mercapto, amino, nitro, C<sub>1-6</sub> alkoxy, C<sub>1-6</sub> alkylthio, and C<sub>1-6</sub> alkylcarboxyl,
  - 15 (c) aromatic and heteroaromatic groups substituted with 2 or 3 adjacent hydroxy groups, and optionally substituted with 1 to 5 non-interfering substituents,



- (d) sugars, optionally substituted with one or more anionic groups selected from sulfate, phosphate, phosphonate, carboxylate, and sulfonate groups,  
(e) peptides and peptide derivatives, and  
(f)  $-C(O)R_3$  and  $-C(O)OR_3$  (where  $R_3$  is selected from the group consisting of (a) through (e) above); and

Y is hydrogen, hydroxy,  $C_{1-6}$  alkoxy, benzyloxy (where the phenyl group is optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl), or  $-OSO_2R_4$  (where  $R_4$  is  $C_{1-6}$  alkyl or phenyl optionally substituted with 1 to 3 substituents selected from halo and  $C_{1-6}$  alkyl);

- and the group of compounds consisting of acacetin, actinorhodine, alizarin, alizarin blue, alizarin orange, alizarinsulfonic acid, alkannin, anthragallol, anthralin, anthrarobin, antharufin, apigenin, apigetrin, apiose, baicalein, baptigenin, 1,2,4-benzenetriol, bostrycoidin, carbidopa, carminic acid, carubicin, cellobiose, centaurein, chloranilic acid, chondrosine, chromotrope 2B, chromotropic acid, chrysamminic acid, chrysarobin, chrysin, chrysophanic acid, cichoriin, citrazinic acid, citromycetin, collinomycin, curvularin, cyanidin, cyanidin 3-glucoside, cyanidin 3-rhamnoglucoside, cyanidin 3,5-diglucoside, cyanidin 3-sophoroside, daphnetin, datiscetin, daunorubicin, delphinidin, deoxyepinephrine, diosmetin, diosmin, dioxethedrine, dopa, dopamine, doxorubicin, droxidopa, echinochrome A, embelin, emodin, ergoflavin, eriodictyol, esculetin, fenoldopam, fomecin A, fomecin B, fraxetin, fraxin, fredericamycin A, fumigatin, fusarubin, fuscine, fustin, galangin, gallein, gallocyanine, gardenin A, gardenin B, gardenin C, gardenin D, gardenin E, genistein, gentisin, granaticin, guamecyclyne, hematein, hydroxysphorobioside, hydroxysphoricoside, icariin, isoquercitrin, kaempferol, kermesic acid, laccaic acid A, laccaic acid B, laccaic acid C, laccaic acid D, leucocyanidin, luteolin, maclurin, menogaril, methylenedigallic acid, morin, oosporein, phenicin, phloroglucide, puberulic acid, puberulonic acid, purpurin, purpurogallin, quercetagenin, quercimritrin, quinalizarin, quinic acid, resistomycin, rhamnetin, rhein, rhodizonic acid, rhodomycin A, rhodomycin B, robinin, ruberythric acid, rufigallol, rutin, scutellarein, tannic acid, tetroquinone, tiron, troxerutin, and tunichrome B1,

but excluding pyrogallol,  
and the pharmaceutically acceptable salts thereof.

36. The drug product of Claim 35 indicated for the treatment of Alzheimer's disease.

5

37. The drug product of Claim 35 indicated for the treatment of Parkinson's disease.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
12 July 2001 (12.07.2001)

PCT

(10) International Publication Number  
**WO 01/49281 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 31/19**,  
31/35, 31/05, 31/70, 31/135, 31/165, 31/325

(21) International Application Number: PCT/US00/35715

(22) International Filing Date:  
28 December 2000 (28.12.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/173,958 30 December 1999 (30.12.1999) US  
09/748,748 26 December 2000 (26.12.2000) US

(71) Applicant: **PROTEOTECH, INC.** [US/US]; 12026 115th  
Avenue N.E., Kirkland, WA 98034 (US).

(72) Inventors: **CASTILLO, Gerardo, M.**; 11725-82nd Ave.  
South, Seattle, WA 98178 (US). **CHOI, Paula, Y.**; Apt.  
108, 11119 Woodenville Drive, Bothell, WA 98011 (US).  
**SNOW, Alan, D.**; 3812 167th Pl. S.W., Lynnwood, WA  
98037 (US).

(74) Agents: **KURZ, Walter et al.**; Heller Ehrman White &  
McAuliffe LLP, 275 Middlefield Road, Menlo Park, CA  
94025-3506 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AT  
(utility model), AU, AZ, BA, BB, BG, BR, BY, BZ, CA,  
CH, CN, CR, CU, CZ, CZ (utility model), DE, DE (utility  
model), DK, DK (utility model), DM, DZ, EE, EE (utility  
model), ES, FI, FI (utility model), GB, GD, GE, GH, GM,  
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,  
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,  
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK  
(utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN,  
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

(88) Date of publication of the international search report:  
10 May 2002

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.



**WO 01/49281 A3**

(54) Title: **POLYHYDROXYLATED AROMATIC COMPOUNDS FOR THE TREATMENT OF AMYLOIDOSIS AND  $\alpha$ -SYNU-  
CLEIN FIBRIL DISEASES**

(57) Abstract: Polyhydroxylated aromatic compounds, and compositions containing them, are useful for the treatment of amyloido-  
sis, especially Alzheimer's disease, and for the treatment of diseases characterized by  $\alpha$ -synuclein fibril formation, especially Lewy  
body disease and Parkinson's disease.

# INTERNATIONAL SEARCH REPORT

International Application No

PC./US 00/35715

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/19 A61K31/35 A61K31/05 A61K31/70 A61K31/135  
A61K31/165 A61K31/235

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP 01 151514 A (NICHIREI CORPORATION) 14 June 1989 (1989-06-14)  abstract  ---  -/--	1-3,5-7, 9,10,17, 19-21, 23-25, 27-29, 31-33, 35,36

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

16 November 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Helps, I

## INTERNATIONAL SEARCH REPORT

International Application No

PC., US 00/35715

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	C. HERTEL ET. AL.: "Inhibition of the electrostatic interaction between beta-amyloid peptide and membranes prevents beta-amyloid toxicity." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE USA, vol. 94, no. 8, August 1997 (1997-08), pages 9412-6, XP001026042 page 9414, column 2, last paragraph to page 9415, column 1, first paragraph page 9413, column 2, paragraph 3 ---	1-3,5-7, 9,10,17, 19-21, 23-25, 27-29, 31-33, 35,36
X	H. NIWANO ET. AL.: "Inhibitory action of amyloid precursor protein against human Hageman Factor (Factor XII)." JOURNAL OF LABORATORY AND CLINICAL MEDICINE, vol. 125, no. 2, 1995, pages 251-6, XP000933861 page 253, column 2 -page 254, column 1 ---	1-3,5, 13,17, 19-21, 23, 27-29, 31,35,36
A	WO 98 24646 A (UNIVERSITY OF HERTFORDSHIRE) 13 August 1998 (1998-08-13)  claims; examples ---	1-3, 5-21, 23-29, 31-36
A	WO 98 09653 A (S.C.R.A.S.) 12 March 1998 (1998-03-12)  claims; examples ---	1-3, 5-21,23, 29,31-36
A	US 5 767 126 A (MARCHBANKS) 16 June 1998 (1998-06-16)  claims; examples -----	1-3, 5-21, 23-29, 31-36

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US 00/35715

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Although claims 1-18 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
2, 3, 20, 21, 28, 29 (complete), 1, 5-19, 23-27, 31-35 (partial)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 2,3,20,21,28,29 (complete) 1,5-19,23-27,  
31-35 (partial)

Use of compounds defined in claim 1 for the treatment of amyloidosis.

2. Claims: 4,22,30,37 (complete) 1,5-19, 23-27,  
31-35 (partial)

Use of compounds of claim 1 for the treatment of disorders caused by alpha-synuclein fibril formation.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PL., US 00/35715

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 01151514	A	14-06-1989	JP 2587842 B2	05-03-1997
WO 9824646	A	11-06-1998	US 5746441 A	05-05-1998
			AU 3822997 A	29-06-1998
			WO 9824646 A1	11-06-1998
			US 5908198 A	01-06-1999
WO 9809653	A	12-03-1998	FR 2753098 A1	13-03-1998
			AU 734296 B2	07-06-2001
			AU 4211197 A	26-03-1998
			EP 0939654 A1	08-09-1999
			WO 9809653 A1	12-03-1998
			JP 2000517336 T	26-12-2000
			NO 991100 A	05-05-1999
			US 6297281 B1	02-10-2001
US 5767126	A	16-06-1998	GB 2264707 A	08-09-1993
			AU 1674492 A	12-01-1993
			EP 0589900 A1	06-04-1994
			WO 9222534 A1	23-12-1992
			JP 6508822 T	06-10-1994